

Minisuperspace, WKB, Quantum States of General Relativistic Extended Objects

S. Ansoldi

*Dipartimento di Matematica e Informatica, Università di Udine, and I.N.F.N. Sezione di Trieste
via delle Scienze, 206 - I-33100 Udine (UD) - ITALY*

Abstract. The dynamics of relativistic thin shells is a recurrent topic in the literature about the classical theory of gravitating systems and the still ongoing attempts to obtain a coherent description of their quantum behavior. Certainly, a good reason to make this system a preferred one for many models is the clear, synthetic description given by Israel junction conditions. Using some results from an effective Lagrangian approach to the dynamics of spherically symmetric shells, we show a general way to obtain WKB states for the system; a simple example is also analyzed.

The study of the dynamics of an (infinitesimally)¹ thin surface layer separating two domains of spacetime is an interesting problem in General Relativity. The system can be described in a very concise and geometrically flavored way using Israel's junction conditions [1, 2, 3]. Starting from these footholds many authors have then tackled the problem of finding some hints about the properties of the *still undiscovered* quantum theory of gravitational phenomena using shells as convenient models. In this context, just as examples of what can be found in the literature, we quote the seminal works of Berezin [5] and Visser [6], that date back to the early nineties, or the more recent [7, 8] and references therein.

What we are going to shortly discuss in the present contribution is set in this last perspective and suggests a semiclassical approach to define WKB quantum states for spherically symmetric shells. This method has already been used in [4].

Let us then consider a spherical shell (we refer the reader to [3] for very concise/clear background material and for definitions). For our purpose the relevant result is equation (4) in [3], i.e. the junction conditions² $K_{ij}^- - K_{ij}^+ \propto S_{ij} - g_{ij}S/2$. K_{ij} is the extrinsic curvature of the shell and can have different values on the two sides (+ and – spacetime regions) of it. S_{ij} is the stress energy tensor describing the energy/matter content of the shell (S is its trace). For a spherical shell these equations reduce to the single condition

$$\varepsilon_-(\dot{R}^2 + f_-(R))^{1/2} - \varepsilon_+(\dot{R}^2 + f_+(R))^{1/2} = M(R)/R, \quad (1)$$

where $f_{\pm}(r)$ are the metric functions in the static coordinate systems adapted to the

¹ Far from being only idealizations of more realistic situations, shells have been extensively used to build concrete models of many astrophysical and cosmological scenarios (for a detailed bibliography, please, see the references in [4]).

² Conventions are as in [9] and the definition of the (quantities relevant to the concept of) extrinsic curvature can be found in [3, 9].




FIGURE 1. Plot of some WKB levels for the example discussed in the text. The curves $a, b, c, d, e, f, g, h, i, j$ correspond to the quantum numbers 1, 2, 3, 4, 5, 10, 20, 30, 40, 50, respectively.

spherical symmetry of the 4-dimensional spacetime regions joined across the shell; $\varepsilon_{\pm} = +1, -1$ are signs and R and $M(R)$ are the the radius (a function of the proper time τ of a co-moving observer³) and the matter content (what remains of S_{ij}) of the spherical shell, respectively. As shown for example in [10, 11, 12] the above equation can be obtained from an effective Lagrangian⁴ $L_{\text{EFF}}(R, \dot{R})$, as a first integral of the second order Euler-Lagrange equation. From $L_{\text{EFF}}(R, \dot{R})$ the momentum conjugated to the single degree of freedom R can be obtained as usual, $P(R, \dot{R}) = \partial L_{\text{EFF}}(R, \dot{R}) / (\partial \dot{R})$. We are not interested in the explicit form of P here, we just point out that it is a function of R and $\dot{R}$ highly *non-linear* in $\dot{R}$. This non-linearity spoils the natural and simple interpretation of the momentum than we know from classical analytical mechanics. Nevertheless we can still solve (1) for the classically allowed trajectories of the shell, using a *standard* analogy with the motion of a *unitary mass particle with zero energy in an effective potential* [16, 17, 18]. This gives $\dot{R}$ as a function of R and, substituting this expression for $\dot{R}$ in $P(R, \dot{R})$, we obtain the *conjugated momentum on a solution of the classical equations of motion*. In what follows we are going to indicate the momentum *evaluated on a classical trajectory* as $P(R; \mathcal{S})$: this emphasizes that it is a function of R , that it depends on the set $\mathcal{S}$ of the other parameters of the problem, but, of course, it is not a function of $\dot{R}$.

By integrating the expression for $P(R; \mathcal{S})$ on a classically allowed trajectory with turning points R_{MIN} and R_{MAX} , we can compute the value of the classical action for that trajectory. It is a function of the set of parameters characterizing the matter content and

³ As usual an overdot, “ $\dot{}$ ”, denotes a total derivative with respect to τ .

⁴ For a more general and deeper discussion see [13] and references therein or also, in addition, [14, 15].

of the geometry, $\mathcal{S}$, and WKB quantum states of the system can now be defined as states for which the above action is a multiple of the quantum

$$S(\mathcal{S}) = 2 \int_{R_{\text{MIN}}}^{R_{\text{MAX}}} P(R; \mathcal{S}) dR \sim n, \quad n = 0, 1, 2, 3, \dots \quad (2)$$

In quantum gravity we expect to have a theory that selects some geometries from the set of all possible ones consistently with quantum dynamics. In our discussion we have limited the treatment to a *minisuperspace approximation*, but, indeed, we see that the quantization condition (2) does select only some of the possible geometries, those in which the parameters of the model are related by the quantum number n as in (2). We can see this more explicitly in the following simple model: a dust shell ($M(R) = m$) separating two domains of anti-de Sitter spacetime with the same cosmological constant ($f_-(r) = f_+(r) = f(r) = 1 + r^2/H^2$): it is possible to prove that a non-trivial junction of the two spacetimes can be performed, although we are not going to discuss this aspect here nor we are going to describe the resulting global spacetime structure. In this case the set of parameters is $\mathcal{S} = \{m, H\}$ and the quantization condition (2) becomes $S(m, H) \sim n$, $n = 0, 1, 2, 3, \dots$. Some levels are plotted in figure 1 and clearly show that given one of the parameters (say m) only a discrete set of values for the other H is allowed: thus the quantization condition restricts the possible values that can be given to the parameters that characterize the spacetime geometry and/or the matter content of the shell. This is consistent with the general picture of a quantum gravitational scenario.

Other applications of this general approach are presently under study. In particular, generalizations to higher dimensions [19] could be relevant, for example, in the context of brane cosmological models.

REFERENCES

1. Israel, W., *Nuovo Cimento B*, **44**, 1 (1966).
2. Israel, W., *Nuovo Cimento B*, **48**, 463 (1967).
3. Barrabes, C., and Israel, W., *Phys. Rev. D*, **43**, 1129–1142 (1991).
4. Ansoldi, S., *Class. Quantum Grav.*, **19**, 6321–6344 (2002).
5. Berezin, V. A., *Phys. Lett. B*, **241**, 194–200 (1990).
6. Visser, M., *Phys. Rev. D*, **43**, 402–409 (1991).
7. Berezin, V., *J. Mod. Phys. A*, **17**, 979–988 (2002).
8. Corichi, A., Cruz-Pacheco, G., Minzoni, A., Padilla, P., Rosenbaum, M., Ryan Jr., M. P., Smyth, N. F., and Vukasinac, T., *Phys. Rev. D*, **65**, 064006(1–13) (2002).
9. Misner, C. W., Thorne, K. S., and Wheeler, J. A., *"Gravitation"*, W. H. Freeman and Company, 1970.
10. Farhi, E., Guth, A. H., and Guven, J., *Nucl. Phys. B*, **339**, 417–490 (1990).
11. Ansoldi, S., Aurilia, A., Balbinot, R., and Spallucci, E., *Class. Quantum Grav.*, **14**, 2727–2755 (1997).
12. Ansoldi, S., *Graduation Thesis (in Italian)*, University of Trieste (1994).
13. Hajicek, P., and Kijowski, J., *Phys. Rev. D*, **62**, 044025(1–5) (2000).
14. Gladush, V. D., *J. Math. Phys.*, **42**, 2590–2610 (2001).
15. Sindoni, L., *Graduation Thesis (in Italian)*, University of Trieste (2004).
16. Berezin, V. A., Kuzmin, V. A., and Tkachev, I. I., *Phys. Rev. D*, **36**, 2919–2944 (1987).
17. Blau, S. K., Guendelman, E. I., and Guth, A. H., *Phys. Rev. D*, **35**, 1747–1766 (1987).
18. Aurilia, A., Palmer, M., and Spallucci, E., *Phys. Rev. D*, **40**, 2511–2518 (1989).
19. Ansoldi, S., Guendelman, E. I., and Ishihara, H. (In preparation).